This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



(51) International Patent Classification 6:		(11) International Publication Number:	WO 99/35917
A23B 7/08, 7/16, A23L 1/09, 1/29	A1	(43) International Publication Date:	22 July 1999 (22.07.99)
(21) International Application Number: PCT/US (22) International Filing Date: 14 January 1999 ((30) Priority Data: 60/071,081 15 January 1998 (15.01.98) (71)(72) Applicant and Inventor: HIRSCHBERG, [US/US]; 25 Tamarack Drive, Hillsborough, C. (US). (74) Agents: WANG, P., Ben et al.; Townsend and T and Crew LLP, 8th floor, Two Embarcadero Cer Francisco, CA 94111–3834 (US).	14.01.9 L Edwa A 9401	BY, CA, CH, CN, CU, CZ, DE, I GE, GH, GM, HR, HU, ID, IL, KR, KZ, LC, LK, LR, LS, LT, I MN, MW, MX, NO, NZ, PL, PI SI, SK, SL, TJ, TM, TR, TT, UA ZW, ARIPO patent (GH, GM, KE ZW), Eurasian patent (AM, AZ, B' TM), European patent (AT, BE, C FR, GB, GR, IE, IT, LU, MC, N (BF, BJ, CF, CG, CI, CM, GA, G SN, TD, TG).	J, AZ, BA, BB, BG, BR, DK, EE, ES, FI, GB, GD, IN, IS, JP, KE, KG, KP, LU, LV, MD, MG, MK, F, RO, RU, SD, SE, SG, UG, US, UZ, VN, YU, E, LS, MW, SD, SZ, UG, Y, KG, KZ, MD, RU, TJ, TH, CY, DE, DK, ES, FI, L, PT, SE), OAPI patent

(54) Title: METHODS OF INFUSING PHYTOCHEMICALS, NUTRACEUTICALS, AND OTHER COMPOSITIONS INTO FOOD **PRODUCTS**

(57) Abstract

This invention provides methods of infusing compositions including phytochemicals, nutraceuticals such as vitamins, herbal extracts, and medicinals into food products, including, e.g., juices, fruits, vegetables, and meats, etc. The resulting infused food products are consumable products which are helpful in alleviating dietary insufficiency and/or to prevent or treat diseases such as cancer, heart disease, Alzheimer's disease, etc.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

United States of America Uzbekistan Vict Nam

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Scnegal
ΑU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
ΑZ	Azerbaijan	GB	United Kingdom	MC	Моласо	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IR	Iteland	MN	Mongolia	UA.	Ukraine
BR	Brazil	lL.	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of Ameri
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Vict Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	zw	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand	LII	Zimozowe
CM	Cameroon		Republic of Korea	PL.	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ.	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SR	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

. . .

20

25

5 METHODS OF INFUSING PHYTOCHEMICALS, NUTRACEUTICALS, AND OTHER COMPOSITIONS INTO FOOD PRODUCTS

CROSS REFERENCE TO RELATED APPLICATIONS

This application is a continuation-in-part of US Serial No.:
60/071,081 filed 01/15/98, entitled "Methods of Infusing Phytochemicals into Food Products" by Edward Hirschberg, and claims priority thereto. The 60/071,081 application is incorporated herein by reference.

FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT Not applicable.

FIELD OF THE INVENTION

This invention provides methods of infusing compositions including phytochemicals, nutraceuticals such as vitamins, herbal extracts, and medicinals into food products, including, e.g., juices, fruits, vegetables, and meats, etc. The resulting infused food products are consumable products which are helpful in alleviating dietary insufficiency and/or to prevent or treat diseases such as cancer, heart disease, Alzheimer's disease, etc.

BACKGROUND OF THE INVENTION

Fruits, vegetables, vitamins, minerals, and phytochemicals included in a diet can enhance disease resistance. Food products such as fruit and vegetables commonly have moisture reduced through various techniques

such as freeze drying, vacuum drying, air drying, osmotic dehydration, etc. to enhance storage, consumption, and shippability.

It has been noted in the conditioning and preservation of food products in their natural state that all food products contain soluble materials which may be lost or altered by contact with solutions of many types. For example, if meats, vegetables, or fruits are placed in water, the water has a tendency to dissolve out the soluble components of these food products, thereby altering their appearance and flavor.

There have been previous attempts to modify food products with useful ingredients. Some of the previous work involved mixing vitamin extracts 10 into a fresh fruit mass, e.g., Elliot Ritchie Alexander, "Vitamin Fruit Composition," US Pat. No. 1,886,931, and Clarence Birdseye, "Process of Improving and Preceiving Food Products," US Per H. 2,419,877. Other involved adding flavors into frozen foodstuffs, such as those disclosed in Harry A. Noyes, "Process of Conditioning Foods," US Pat. No. 2,317,397. More 15 recent work described using osmosis to dehydrate and preserve foods, e.g. Brimelow and Brittain, "Contiguous Food Impregnation," US Pat. No. 4,183,963, and Peter Butland, "Process for Producing a Natural Fruit Candy," US Pat. No. 4,355,050. There have also been efforts made to infuse certain flavors into foods using osmosis, such as those disclosed in Lewis and Lewis, 20 "Dehydrated Vegetables," US Pat. No. 4,683,141, Karim Nafisi-Novaghar, "Method of Making Dried Fruits," US Pat. No. 5,000,972, and Aebi, et al., "Process for Preparing Dehydrated Vegetable Products," US Pat. No. 5,368,873, and using a degradation enzyme, e.g. as in Poulose and Boston, "Enzyme Assisted Degradation of Surface Membranes of Harvested Fruits and 25 Vegetables," US Pat. No. 5,037,662. Some have also explored methods of preserving fruits and vegetables using deep freezing, e.g., Berey et al., "Fruit and Vegetable Preservation Using Deep Freeze - by Complete or Partial Withdrawal of Free Water and Partial Replacement of it with Additives,

15

20

Improving Appearance, Flavour and Shelf-life," e.g., in Hungarian patent HU 47823.

However, none of these prior art methods are satisfactory in terms of their ability to infuse nutrients into food products in general. The reason is that the above processes result in a loss of phytochemicals, vitamins and minerals from the food, resulting in a corresponding loss of nutritional value and decreasing the disease-preventative value of the food. A need exists in the art to provide improved methods of food preparation and storage, as well as to provide efficient mechanisms for infusing osmotically dried fruits, vegetables, and other food products. This invention fills these and other needs.

SUMMARY OF THE INVENTION

techniques such as freeze drying, vacuum drying, air drying, osmotic dehydration, etc. One process described herein uses osmotic dehydration, in which higher osmotic pressure outside a food particle causes moisture to migrate out of the food. This is done with sugars or solutions of sugars, e.g., high fructose corn syrup, or others. During the migration time, solutes in the solutions fill voids left by the water that was removed by infusion. In the invention, flavors, colors, vitamins, minerals, and phytochemicals such as isoflavonals, lycopene, resveratol, indocarbonals, anthocyanins, etc., as well as, e.g., soluble fiber, are infused along with the osmotic dehydration solute by adding these desired additives in controlled dosage to the osmotic dehydration solution.

The invention provides a method of infusing a composition into a food product by increasing the brix of an osmotic dehydration solution containing the food product, over a period of time. The food product is incubated with the composition, thereby infusing the composition into the food

15

20

25

product. These steps are optionally performed separately, sequentially, or simultaneously.

Typically, the composition used in the method is a phytochemical, a nutraceutical, or other desirable food additive. In one significant aspect of the invention, the food product is predried before the composition is infused into the food product. Alternatively, the osmotic dehydration solution optionally includes the composition to be infused into the food product.

The brix of the osmotic dehydration solution is typically varied from about 20 °B to about 80 °B. The brix of the osmotic dehydration solution is optionally changed daily, e.g., for a period of at least about one week. The method of the present invention can be performed at various temperatures such as about room temperature or up to about 50 °C, e.g., where the brix of the osmotic dehydration solution is about 77 °B. The infection method fartime comprises stirring or circulating the osmotic dehydration solution, thereby increasing the rate of infusion of the composition into the food product. In one example, the infusion method is performed where the food product includes a strawberry and the brix of the osmotic dehydration solution is about 40 °B. In another example, the food product can include a marionberry and the osmotic dehydration solution includes high fructose corn syrup (HFCS). In another example, the food product used in the present invention includes a sliced cranberry, and the osmotic dehydration solution includes a phytochemical and HFCS, and the brix of the osmotic dehydration solution is about 77 °B.

In another aspect of the invention, when the food product includes fresh sliced carrot, and the brix of the osmotic dehydration solution is about 77 °B, the infusion method further includes heating the osmotic dehydration solution containing HFCS, Saw Palmetto, and Gingko Biloba, and incubating the solution overnight. Typically, the osmotic dehydration solution includes a dehydration solute such as high fructose corn syrup, dextrin, starch, gelatin, pectin, juice concentrate, or soy isolate. Optionally, the food product can be

10

15

25

coated with a coating substance such as gelatin, pectin, or starch to form a coated food product, and the phytochemical is infused into the coated food product.

The dehydration solution is optionally heated, and/or continuously circulated. The solution is typically circulated through a tube or a pipe, or passed over a perforated conveyor in either a continuous or discontinuous process. In yet another aspect of the invention, the food product is freeze dried to 10% or lower residual moisture before infusion occurs.

Compositions to be infused include can be phytochemicals, or nutraceuticals such as: vitamins, minerals, isoflavoronals, lycopene, resveratol, indocarbonals, anthocyanins, soluble fiber, high protein rice, soy isolate or others. Optionally, the composition to be infused can include a flavor, or a color. The food product is optionally a vagatille such that correct, or built pepper, e.g., where the osmotic dehydration solution includes a low dextrose and L-Carnitine.

In a further aspect of the invention, excess water is removed by drying a mixture of the food product and the composition after incubation of the food product with the composition. The food product includes, e.g., fruit, fruit juice, a vegetable, vegetable juice, ground liver, chicken, salmon, or the like. For example, the food product can be an apple, a carrot, or a pet treat. The 20 composition used in the method of this invention can be a medicinal capable of providing a medical or dietary benefit to a human or an animal. For example, in one embodiment, prunes or prune juices are infused with Chinese herbal medicine to form a mixture for treatment or prevention of irritable bowel syndrome.

In one aspect of the invention, where the food product is a fruit juice or vegetable juice, and the composition is a phytochemical, the phytochemical is infused into the fruit juice or vegetable juice and further formed into a solid or semisolid mixture.

15

20

25

The mixture thus formed can be made into a trail mix, e.g., with at least two kinds of food products infused with phytochemicals. In a preferred embodiment, the mixture contains high protein rice, or soy isolate. In another preferred embodiment, the mixture is formed into a firm mass of material mixed with pectin or gelatin. Alternatively, the mixture is coated with a coating substance such as gelatin, pectin, or starch.

In a further aspect of the invention, a flavoring is infused into a fruit using the following steps: (a) soaking the fruit in a solution of potassium sorbite, calcium lactate, citric acid, glycerol, and the flavoring for at least three days; (b) removing 20% of the solution by weight each successive day; (c) replacing the removed solution with about 77 °B brix HFCS until the brix of the solution reaches about 65 °B; (d) rinsing the solution off the fruit or vegetable; and (e) drying the first or vegetable. The first principal lay includes, e.g., blueberry, strawberry, marionberry, cranberry, or the like. Optionally, anthocyanins, vitamin C, or other compounds are added to the solution.

DEFINITIONS

As used herein, the following terms have the meanings ascribed to them unless specified otherwise. All other terms have their meanings as commonly understood by one skilled in the art.

The term "food product" refers to material of either plant or animal origin, or of synthetic sources, that contain an essential body nutrient such as a carbohydrate, protein, fat, vitamin, mineral, etc. Examples include meats, fruits, vegetables, grains, nuts, and the like.

The term "phytochemical" refers to dietary compounds or chemicals derived from plants which provide a health benefit when consumed by humans or animals. Examples of such compounds or chemicals are lycopene, isoflavones, etc.

20

The term "nutraceutical" refers any compounds or chemicals that can provide dietary or health benefits when consumed by humans or animals. Examples of nutraceuticals include vitamins, minerals, and others.

The term "osmotic dehydration" refers to the process which occurs when food products are submerged in a solution having higher concentration of solutes outside of the food product than inside of it, and when water is caused to migrate from within the food product to outside the food product, thereby causing the food product to be dehydrated.

The term "brix" refers to a unit measurement for measuring the

density of the concentration of a chemical such as sugar in a solution. As used
in the art, brix refers to a concentration in percent of sugar by weight according
to the Brix scale. Brix scale here refers to a hydrometer scale for sugar
solutions so graduated that its readings in degrees Prince or confined
temperature represent percentages by weight of sugar in a solution. See

Webster's Third New International Dictionary, unabridged, published by G. &
C. Merriam Company, Springfield, Massachusetts.

The term "pretreating" refers to a process where a food product is treated chemically or physically before it is infused with a composition.

The term "predried" refers to a process where a food product is dehydrated by various means before it is infused with a composition. For example, the means of dehydration can be by freeze drying such that the food product is in freeze dried format when it is incubated with the composition.

The term "HFCS" refers to high fructose corn syrup.

The term "pet treat" refers to a food product consumed by a pet or
livestock (i.e. a non-human, domesticated animal), such as a dog or cat biscuit,
or other dog or cat treat. Examples of pets includes dogs, cats, birds, snakes,
turtles, frogs, toads, rabbits, rats, mice, gerbils, hamsters, fish, and the like.
Examples of livestock include pigs, cattle, cows, horses, chickens, ducks, fish,

and other animals. Animal treats can, of course, also be produced for undomesticated wild animals.

The term "herbal medicine" refers to naturally occurring compositions such as Chinese herbal medicine that are used by Traditional Chinese Medicine practitioners in curing diseases. Such compounds are generally derived from plant sources. Examples of herbal medicine include Dang Shen, Hua Xiang, Fang Feng, Yin Chen, Fu Ling, Bai Zhi, Bai Shao, Mu Xiang, Wu Wei Zi, etc.

The term "coating substance" refers to a composition that can be applied to the food product to be infused a phytochemical or other composition to form a protective membrane or coating or layer over the food product.

Examples of such coating substance are gelatin, pectin, and starch.

The term "trail mix" refere to a food product containing c. Len two kinds of fruits, vegetables, nuts, or other foods. In the methods of the invention, at least one of the at least two kinds of food products is infused with a composition such as phytochemical, or nutraceutical.

DESCRIPTION OF THE PREFERRED EMBODIMENT

The present invention relates to processes to infuse

20 phytochemicals, nutraceuticals, and other compositions into food products, so as
to form mixtures that can be consumed by humans or animals. The method
disclosed herein can be used to produce snacks or other food items containing
vitamins, minerals, other nutrients or medicinals so as to provide dietary
benefits to humans or animals.

25

10

15

INFUSION CONDITIONS

The infusion process is conducted by gradually increasing the brix of the osmotic dehydration solution from 20°B to 65°B, usually daily, typically over a one to two week period. Optionally, the infusion is speeded up by the

15

20

25

use of heat, i.e., from ranges of 20-80 °C, optionally 30-70 °C, and generally 40-60 °C, and preferably at about 50 °C, and a high strength of solution, e.g., 77°B, to under 24 hours. Stirring of the solution is optionally used to speed the infusion. The brix of the osmotic dehydration solution is typically varied from about 20 °B to about 80 °B. Preferably, the brix can be from 30-70 °B, or from 40-60 °B, or from 60-80 °B. The brix of the osmotic dehydration solution is optionally changed daily, e.g., for a period of at least about one week. It is expected that one of skill in the art is fully able to adjust the brix of a solution by adding solute to the solution, and/or by partial or complete replacement of the solution.

The method of the present invention can be performed at various temperatures, such as at about room temperature or up to about 50 °C to thin out the delaydration solute for greater absorption. Optionally, the blik of the osmotic dehydration solution is from 20-80 °B, depending on the desired condition of infusion. For example, in one embodiment, infusion is conducted at about 77 °B.

For example, the infusion method can be performed when the food product includes a strawberry with the brix of the osmotic dehydration solution being about 40 °B. In another embodiment, the food product can include a marionberry and the osmotic dehydration solution includes high fructose corn syrup (HFCS). Optionally, the food product used in the present invention can include a sliced cranberry, and the osmotic dehydration solution includes a phytochemical and HFCS, and the brix of the osmotic dehydration solution is about 77 °B.

The osmotic dehydration solution includes a sugar component such as high fructose corn syrup. Typically, it includes at least about 35% to about 100% fructose, and preferably about 42% to about 90% fructose. The balance of the sugar may be comprised of dextrose or any of a number of saccharide materials including monosaccharides, disaccharides and polysaccharides and

10

15

their degradation products, e.g., pentoses including aldopentoses, ketopentoses like xylose and arabinose, a deoxyaldose like rhamnose, hexoses and reducing saccharides such as aldohexoses like glucose, galactose and mannose; the ketohexuloses, like sorbose and xylulose; disaccharides, like maltulose, lactose and maltose; nonreducing disaccharides such as a sucrose, other polysaccharides such as dextrin and raffinose, and hydrolyzed starches which contain as their constituents oligosaccharides. The balance of the sugar may be of a low molecular weight so as to offer a substantial effect in increasing the osmotic pressure of the sugar solution. The balance of the sugar may also be comprised of polyhydric alcohols such as glycerol and the like. When polyhydric alcohols are employed, they preferably comprise only about 1 to about 10% of the sugar component.

The infusion can be perfermed in various constitute teachly available in the art. For example, the containers for the infusion are typically made of plastic or other polymer materials. Other materials that can be used for the container include stainless steel, aluminum, glass, or other ceramics. When applicable, the heat source can be applied to the container internally or externally, or both, by a heating means such as a heating rod, open burner, closed burner, electrical heating element, or the like.

20

25

PHYTOCHEMICALS USED IN INFUSION

The osmotic dehydration solution optionally includes any of a variety of vitamins, including vitamin A (e.g., vitamin A_1 , retinol, axerophthol, α -carotene, β -carotene, γ -carotene), B vitamins (e.g., B_1 vitamins including: thiamin, aneurin, thiamine, pyrophosphate, cocarboxylase; B_2 vitamins including riboflavin, vitamin G, lactoflavin, hepatoflavin, ovoflavin, verdoflavin, riboflavin mononucleotide, FMN, riboflavin dinucleotide, FAD), Vitamin C (ascorbic acid, antiscorbutic vitamin, dehydroascorbic acid), D

15

20

25

WO 99/35917 PCT/US99/00181

11

(antirachitic vitamin, vitamin D₂, D₃, cholecalciferol, etc), Vitamin E, Vitamin K, and the like.

For an overview of vitamins, minerals and biologicals, and their sources and formulations, *see*, Kirk-Othmer (1984) Encyclopedia of Chemical Technology, 3rd edition, volume 24 A, Wiley-Interscience Publication, John Wiley & Sons, New York, and Kirk-Othmer (1994) Encyclopedia of Chemical Technology, 4th edition, volume 11, A Wiley-Interscience Publication, John Wiley & Sons, New York. For additional reference materials, *see also*, The Merck Index (1989), An Encyclopedia of Chemicals, Drugs, and Biologicals, 11th Edition, Published by Merck & Co., Inc., Rahway, NJ (or any later versions of the Merck Index).

The solution optionally includes flavoring compounds or other feed additives (e.g., colors and coloring agents, antioxidents, measuratives, flavor enhancers, enzymes, emulsifiers, firming agents, formulation aids, nonnutritive sweeteners, processing aids, stabilizers, etc.) such as those described Kirk-Othmer (1994) Encyclopedia of Chemical Technology, 4th edition, volume 11, *supra* or any other available source. Similarly, the solution can include oils, solubilizers and the like to aid in solubilizing components which have poor solubility in aqueous solutions. After infusion, the food product is optionally further dried for stability and convenience of use. Other food processing techniques are also applicable and described *id*. Some foods that have low structural integrity or need a skin to transfer them are coated with an edible polymer. This can include liquids that have been solidified with gums, pectins, etc., and are thereafter coated with edible polymers.

In one aspect, the present invention provides for fruits and vegetables with enhanced properties such as higher vitamin content, soluble fiber, amino acids, or the like. Using the methods herein, it is possible to add fiber, vitamins, flavorings, preservatives, and the like, to most fruits and vegetables, as well as to many other foods. In one embodiment, it is desirable

10

20

25

to fortify a fruit or vegetable with a vitamin naturally present in the vegetable, thus giving it superior nutritional value. For instance, a carrot can be fortified with vitamin A. In other embodiments, it is desirable to fortify the fruit or vegetable with vitamins, minerals, antioxidants, fiber or the like, which are not naturally present in the fruit or vegetable.

In another aspect, herbal or other natural products with beneficial qualities such as ginseng, taro root extract, other Chinese herbal medicine, or the like can be added to the fruit. Similarly, natural hormones such as melatonin can be added to provide various therapeutic benefits (e.g., to combat insomnia), as can pharmaceuticals. One of skill will appreciate that there are many formulations of vitamins, medicinals, herbal medications, natural hormones, etc., which are broadly available, which can be added to the food products by the methods of the invention.

15 FREEZE DRYING BEFORE INFUSION

All food products contain moisture. The process of osmotic dehydration and infusion are limited by the differential in osmotic pressure from outside the food product and a lesser pressure from within the food product. The process of overcoming the inner pressure is difficult and time consuming and never completely replaces all of the residual moisture in the food product.

The process described here starts by freeze drying the food product to under 10% residual moisture. The dry product can be thought of as a dry sponge with voids where moisture previously was and which has little resistance from absorbing media.

One application of the infusion method includes placing the food product in a container and adding high fructose corn syrup. The syrup essentially fills the voids left as the moisture is removed. The process can be as short as minutes to 24 hours, e.g., until the syrup has been substantially

absorbed. The product is removed, rinsed of surface syrup (e.g., in water) and vacuum dried to the desired moisture level.

Ways that the absorption can be significantly speeded up include using a hot syrup, or by heating the container in which the absorption takes place, in order to thin out the syrup. Additionally, circulating the syrup effectively improves absorption due to dynamic contact of the food product with the syrup. Since food products are filled with air which replacing the moisture from within, they tend to float and must be held submerged. Some of the food products can become crushed or deformed, so care is taken to prevent damage during processing.

MIXING AND CIRCULATING THE INFUSION SOLUTION

An additional process of continuously moving both the food product and syrup has been devised. One process includes pumping both the food product and syrup (or other osmotic dehydration solution) through tubes or pipes. Alternatively, both can be mechanically conveyed, or the food product can be conveyed through the osmotic dehydration solution. For example, the food product can be passed over a perforated conveyer through the osmotic dehydration solution. The food product exits the solution on the conveyor and can subsequently be rinsed, oiled, placed in trays and redried. While freeze drying facilitates the osmotic dehydration and infusion, alternative media can be used other than high fructose corn syrup, including dextrin, starch, gelatin, pectin, juice concentrate, etc. The ability to use solutions other than HFCS is a particular benefit of the process outlined herein.

25

10

15

20

APPLYING THE INFUSION TO VARIOUS FOOD PRODUCTS

Food products can be formed into a particular shape, e.g., juice concentrate (e.g., from fruit or vegetable homogenate). For example, food products can be shaped into cubes, which can be of any geometric shape, which

10

15

are then mixed with fruit or vegetable powders having phytochemicals. The mixtures are combined with high fructose corn syrup, and are then dried to form a final product consumed as a snack. Alternatively, the above juice concentrate can be mixed with high protein rice, or soy isolate before infusion with phytochemicals, and a different snack containing high protein rice or soy isolate can be manufactured following a similar process.

Another process to infuse phytochemicals into food product is to start with juice drinks containing phytochemicals. Adequate amounts of pectin or gelatin are mixed with the juice drinks so as to form a firm mass or ball. The firm ball can be cut into various shapes, e.g., dices, or cubes. Next, HFCS is incorporated into the juice cubes. Later, the juice cubes are dried and infused with phytochemicals.

Optionally, the cubes are control with placin pootin, resich, or color coating substances prior to or after drying. After the infusion with phytochemicals, or medicinals, the food product is formed into a bar. Such bars of infused food products can provide a strong medical benefit to the body. Depending on the particular medicinal that is being infused into the food product, they can be effective in the prevention of cancer, heart disease, Alzheimer's, and the AIDS disease. For example, anti-oxidants have been implicated in reducing the risk of these diseases.

Another variation of the above is to form a trail mix of food product containing a variety of fruits, each infused with a phytochemical, or medicinal, so as to provide the consumer with a combined medical or dietary benefits to the body.

25

20

INFUSED PET TREATS

To produce pet treats infused phytochemicals, pet foods such as ground liver, ground chicken, and others are formed into cubes. Then infusion is performed with the pet food cubes to infuse medicinals or phytochemicals for WO 99/35917 PCT/US99/00181

pets, or other livestock, or even for wild animals. These pet treats infused with medicinals can be used a medicine to prevent, ameliorate, or cure diseases. These infused treats are more effective than pills because they are more acceptable to the animals. For different pets, different starting food products are used. For example, for cats, ground salmon meats can be used, while for horses, apples and carrots can be infused.

INFUSING VARIOUS FRUITS AND VEGETABLES

5

10

15

25

Specific preferred examples of infusing fruits with various phytochemicals are described below. One of skill will appreciate that a variety of parameters can be altered to achieve essentially similar results.

Blueberries are soaked with equal parts of water and solution containing potassium scribsto, calcium lactato, citric acid, glycerol and flave in for 3 days. Each successive day, 20% of the solution by weight is removed and replaced with 77 °B brix HFCS. This is continued until the brix reaches 65 °B. The blueberry solution is drained, the blueberries rinsed, oiled, and placed into trays and vacuum dried to desired dryness. Anthocyanins and vitamin C are added in early exchanges for enhancement.

Similarly, strawberries are infused under modified conditions from above. For example, the starting solution containing the phytochemical can be 20 about 40 °B brix. In the case of cranberries, the fruits are sliced and freeze dried prior to infusion. Subsequently, the slices are soaked with 77 °B HFCS containing phytochemicals to be infused. In the case of marionberries, the fruits are infused with phytochemicals following similar osmotic dehydration and infusion procedure as described above.

Prunes or prune slices are combined with prune concentrate and infused with Chinese herbal medicine extract such as those disclosed in Bensoussan, et al, "Treatment of Irritable Bowel Syndrome with Chinese Herbal Medicine", JAMA, Vol. 280, No. 18, pp. 1585-1589 (1998). The mixture is

10

15

20

redried, and formed into a snack form such as a bar. Such snacks can function as a laxative, and can be used in the prevention of colon cancer. Alternatively, prune juice can be mixed with the above mentioned Chinese herbal medicine and gelatin, and forming a gel mixture which is subsequently soaked in prune concentrate. The Gelle mixture is then freeze dried, pulverized, and formed into a tablet or a discrete piece.

Many vegetable juices are infused with phytochemicals, nutraceuticals, and other compositions using the infusion methods of this invention. In some cases, the vegetables juices are mixed with gelatin containing one or more nutraceuticals and then freeze dried before further infusion.

Other vegetables such as carrots, bell peppers, cucumbers, tomatoes, eggplants, celery, sprouts, etc. are freeze dried first. They are then soaked with a low dextrose (e.g., Maltodextrose) containing L-Carnitine in order for the infusion to occur. In the case of fresh sliced carrots, they are first soaked in a 77 °B brix HFCS containing Saw Palmetto, Gingko Biloba. Next, they are incubated overnight in a heated solution in order to speed up the infusion.

Many other aspects will be apparent on reading this disclosure. All references cited herein are incorporated by reference in their entirety.

WO 99/35917 PCT/US99/00181

WHAT IS CLAIMED IS:

1	1.	A method of infusing a composition into a	food product, the method
2		comprising:	

- a. increasing brix of an osmotic dehydration solution containing the
 food product over a period of time; and
- b. incubating the food product with the composition, thereby infusing
 the composition into the food product.
- 1 2. The method of claim 1, wherein the composition is a phytochemical.
- 1 3. The method of claim 1, wherein the composition is a nutraceutical.
- 1 4. The method of claim 1, wherein the food product is predried before
- 2 infusing the composition into the food product.
- 1 5. The method of claim 1, wherein the solution comprises the composition.
- 1 6. The method of claim 1, comprising performing step (a) and step (b)
- 2 separately.
- 1 7. The method of claim 1, comprising performing step (a) and step (b)
- 2 simultaneously.
- 1 8. The method of claim 1, comprising varying the brix of the osmotic
- dehydration solution from about 20 °B to about 80 °B.
- 1 9. The method of claim 1, comprising increasing the brix of the osmotic
- dehydration solution daily for a period of at least one week.

WO 99/35917 PCT/US99/00181 18

- The method of claim 1, wherein the composition is a phytochemical or 10. 1
- nutraceutical selected from: vitamins, minerals, isoflavoronals, lycopene, 2
- resveratol, indocarbonals, anthocyanins, soluble fiber, high protein rice, 3
- 4 and soy isolate.
- The method of claim 1, wherein the composition comprise a flavor, or a 1 11. 2 color.
- The method of claim 1, wherein step (a) or step (b) is performed at room 1 12. 2 temperature.
- The method of claim 1, wherein step (a) or step (b) is performed at about 1 13.
- 50 °C, and the brix of the osmotic dehydration solution is about 77 °B. 2
- The method of claim 1, further comprising stirring or circulating the 1 14.
- osmotic dehydration solution, thereby increasing the rate of infusion of 2
- the composition into the food product. 3
- The method of claim 1, wherein the food product comprises a strawberry 1 15.
- 2 and the brix of the osmotic dehydration solution is about 40 °B.
- The method of claim 1, wherein the food product comprises a 1 16.
- marionberry and the osmotic dehydration solution contains high fructose 2
- com syrup (HFCS). 3
- The method of claim 1, wherein the food product comprises fresh sliced 1 17.
- carrot, and the brix of the osmotic dehydration solution is about 77 °B, 2
- said method further comprising heating the osmotic dehydration solution 3
- containing HFCS, Saw Palmetto, and Gingko Biloba, and incubating said 4
- 5 solution overnight.

PCT/US99/00181

- 1 18. The method of claim 1, further comprising coating the food product with
- a coating substance comprising gelatin, pectin, or starch to form a coated
- food product, and infusing the phytochemical into the coated food

19

- 4 product.
- 1 19. The method of claim 1, wherein the osmotic dehydration solution
- 2 comprising a dehydration solute selected from: high fructose corn syrup,
- dextrin, starch, gelatin, pectin, juice concentrate, and soy isolate.
- 1 20. The method of claim 19, further comprising heating the dehydration
- 2 solute.
- 1 21. The method of claim 19, further comprising continuously circulating the dehydration solute.
- 1 22. The method of claim 19, further comprising circulating the dehydration
- solute and the food product through a tube or a pipe.
- 1 23. The method of claim 19, comprising passing the dehydration solute and
- the food product over a perforated conveyor, and collecting the food
- 3 product therefrom.
- 1 24. The method of claim 1, further comprising pretreating the food product
- by freeze-drying the food product to 10% or lower residual moisture.
- 1 25. The method of claim 24, wherein the food product comprises a sliced
- 2 cranberry, and the osmotic dehydration solution contains the
- phytochemical and HFCS, and the brix of the osmotic dehydration
- 4 solution is about 77 °B.

WO 99/35917 PCT/US99/00181

- 1 26. The method of claim 24, wherein the food product is a vegetable selected
- 2 from: carrot, and bell pepper, and the osmotic dehydration solution
- 3 comprises a low dextrose and L-Carnitine.
- 1 27. The method of claim 1, said method further comprising removing excess
- water by drying a mixture of the food product and the composition after
- incubation of the food product with the composition in step (b).
- 1 28. The method of claim 27, wherein the food product is selected from: fruit,
- fruit juice, vegetable, vegetable juice, ground liver, chicken, and salmon.
- 1 29. The method of claim 27, wherein the food product comprises apple or
- 2 carrot.
- 1 30. The method of claim 27, wherein the food product comprises a prune or a
- 2 prune slice, and the composition comprises an herbal medicine.
- 1 31. The method of claim 27, wherein the food product comprises prune juice,
- and the composition comprises an herbal medicine.
- 1 32. The method of claim 1, wherein the food product is a pet treat.
- 1 33. The method of claim 1, wherein the composition is a medicinal capable
- of providing a medical or dietary benefit to a human.
- 1 34. The method of claim 1, wherein the composition is a medicinal capable
- of providing a medical or dietary benefit to an animal.
- 1 35. The method of claim 1, wherein the food product is a fruit juice or
- vegetable juice, and the composition is a phytochemical, said method
- further comprising infusing the phytochemical into the fruit juice or
- 4 vegetable juice and forming a solid or semisolid mixture.

WO 99/35917 PCT/US99/00181 21

1	36.	The method of claim 35, further comprising forming a trail mix
2		containing at least two food products infused with phytochemicals.

- The method of claim 35, further comprising admixing the mixture with 37. 1 2 high protein rice.
- The method of claim 35, further comprising admixing the mixture with 38. 1 2 soy isolate.
- The method of claim 35, further comprising adding pectin or gelatin to 39. 1 the mixture, thereby forming a firm mass of material. 2
- The method of claim 39, further comprising coating the firm mass of 1 40. material with a coating substance comprising gelatin, pectin, or starch. 2
- A method of infusing a flavoring into a fruit or a vegetable, the method 1 41. comprising: 2
- soaking the fruit or vegetable in a solution of potassium sorbite, 3 calcium lactate, citric acid, glycerol, and the flavoring for at least 4 5 three days;
- removing 20% of the solution by weight each successive day; 6 b.
- replacing the removed solution with about 77 °B brix HFCS until 7 c. the brix of the solution reaches about 65 °B; 8
- 9 rinsing the solution off the fruit or vegetable; and d.
- 10 e. drying the fruit or vegetable.
- The method of claim 41, wherein the fruit is selected from: blueberry, 42. 1
- 2 strawberry, marionberry, and cranberry.
- The method of claim 41, further comprising adding one or more of: l 43.
- 2 anthocyanins, and vitamin C, to the solution.

INTERNATIONAL SEARCH REPORT

International application No. PCT/US99/00181

IPC(6) US CL According	ASSIFICATION OF SUBJECT MATTER :A23B 7/08, 7/16; A23L 1/09, 1/29 :424/ 439; 426/102, 293, 302, 303, 321, 639, 65 to International Patent Classification (IPC) or to be	0 oth national classification and IPC
	LDS SEARCHED	
	documentation searched (classification system followed)	
	424/ 439; 426/102, 293, 302, 303, 321, 639, 650	
Documents	tion searched other than minimum documentation to	he extent that such documents are included in the fields searched
Electronic	data base consulted during the international search	name of data base and, where practicable, search terms used)
C. DOC	CUMENTS CONSIDERED TO BE RELEVANT	
Category*	Citation of document, with indication, where	appropriate, of the relevant passages Relevant to claim No.
X, P — Y,P	US 5,718,939 A (NUGENT) 17 Febr lines 61-70, col. 4, lines 6-51, col. 5	uary 1998, abstract and col.2, 1-43
x x	US 2,785,071 A (MATHEWS) 12 M US 1,906,295 A (WICKENDEN) 02	
y x	60. US 4,551,348 A (O'MAHONY et al.) and col. 7, lines 45-55 and col. 8, lin US H950 A (O'BRIEN et al.) 06 Aug lines 48-68.	05 November 1985, abstract 1 es 1-50.
A* door to b E* earli L* door cited	cr documents are listed in the continuation of Box (cial categories of cited documents: ument defining the general state of the art which is not considered e of particular relevance ior document published on or after the international filing date ument which may throw doubts on priority claim(s) or which is d to establish the publication date of another citation or other ital reason (as specified)	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be
P* docu	ment referring to an oral disclosure, use, exhibition or other na ment published prior to the international filing date but later than priority date claimed uctual completion of the international search	combined with one or store other such documents, such combined with one or store other such documents, such combination being obvious to a person skilled in the art *&" document member of the same patent family
25 MARCI	H 1999	Date of mailing of the international search report 15 APR 1999
Commission Box PCT Washington, Caesimile No		HELEN PRATI Telephone No. (703) 308-1978

INTERNATIONAL SEARCH REPORT

International application No. PCT/US99/00181

		FC1/0399/001		
C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT				
Category*	Citation of document, with indication, where appropriate, of the relev	ant passages	Relevant to claim No.	
x	US 4,775,545 A (AUGUSTINE et al.) 04 October 1988 and col. 5, lines 15-20.	, abstract	1, 24	
X Y	US 5,510,131 A (POULOSE et al.) 23 APRIL 1996, at col. 1, lines 10-29, and col. 7, lines 35-59.	ostract and	1, 2, 3, 5, 10, 11, 15, 25-40	
			1-3, 5, 10, 11, 15, 25-40	
	US 4,917,910 A (HSIEH et al.) 17 April 1990, abstract and 6.	and col. 5	35, 36-38	